arXiv:1407.4403v1 [math.DG] 16 Jul 2014

LIE GROUPS AS 3-DIMENSIONAL ALMOST CONTACT B-METRIC MANIFOLDS

HRISTO MANEV, DIMITAR MEKEROV

Abstract. The object of investigations are almost contact B-metric structures on 3-dimensional Lie groups considered as smooth manifolds. There are established the existence and some geometric characteristics of these manifolds in all basic classes. An example is given as a support of obtained results.

Introduction

The differential geometry of the almost contact metric manifolds is well studied (e.g. [2]). The beginning of the investigations on the almost contact manifolds with B-metric is given by [5]. These manifolds are the odd-dimensional counterpart of the almost complex manifolds with Norden metric [4, 6], where the almost complex structure acts as an anti-isometry on the metric. Almost contact B-metric manifolds are investigated and studied in [5, 12, 13, 14, 15, 16, 17, 19].

An object of special interest is the case of the lowest dimension of the considered manifolds. Curvature properties of the 3-dimensional manifolds of this type are studied in [11, 18, 20].

The basic problem considered in this work is the existence and the geometric characteristics of almost contact B-metric structures on 3-dimensional Lie algebras. The main result is a construction of the considered structures and their characterization in all basic classes of the almost contact B-metric manifolds in the classification given in [5].

The present paper is organized as follows. In Sect. 1 we recall some facts about the almost contact B-metric manifolds. In Sect. 2 we construct different types of contact structures on Lie algebras in dimension 3. In Sect. 3 we consider an example in relation with the investigations in the previous section.

1. Preliminaries

1.1. **Almost contact manifolds with B-metric.** Let $(M, \varphi, \xi, \eta, g)$ be an almost contact manifold with B-metric or an *almost contact B-metric manifold*, i.e. M is a $(2n+1)$-dimensional differentiable manifold with an almost contact structure (φ, ξ, η) consisting of an endomorphism φ of the tangent bundle, a vector field ξ,

1991 *Mathematics Subject Classification.* 53C15, 53C50, 53D15.

Key words and phrases. Almost contact structure, B-metric, Lie group, Lie algebra, indefinite metric.

its dual 1-form η as well as M is equipped with a pseudo-Riemannian metric g of signature $(n+1,n)$, such that the following algebraic relations are satisfied [5]:

$$\begin{aligned} &\varphi\xi=0, \qquad \varphi^2=-\operatorname{Id}+\eta\otimes\xi, \qquad \eta\circ\varphi=0, \qquad \eta(\xi)=1, \\ &g(\varphi x,\varphi y)=-g(x,y)+\eta(x)\eta(y), \end{aligned} \tag{1}$$

where Id is the identity and x, y are elements of the algebra $\mathfrak{X}(M)$ of the smooth vector fields on M.

Further x, y, z will stand for arbitrary elements of $\mathfrak{X}(M)$ or vectors in the tangent space T_pM of M at an arbitrary point p in M.

The associated metric $\tilde{g}$ of g on M is defined by $\tilde{g}(x,y)=g(x,\varphi y)+\eta(x)\eta(y)$. The manifold $(M,\varphi,\xi,\eta,\tilde{g})$ is also an almost contact B-metric manifold. The signature of both metrics g and $\tilde{g}$ is necessarily $(n+1,n)$. By ∇ and $\tilde{\nabla}$ we denote the Levi-Civita connection of g and $\tilde{g}$, respectively.

A classification of almost contact B-metric manifolds is given in [5]. This classification, consisting of eleven basic classes $\mathcal{F}_1$, $\mathcal{F}_2$, $\ldots$, $\mathcal{F}_{11}$, is made with respect to the tensor F of type (0,3) defined by

$$F(x,y,z)=g\big((\nabla_x\varphi)\,y,z\big).$$

The following properties are valid:

$$F(x,y,z)=F(x,z,y)=F(x,\varphi y,\varphi z)+\eta(y)F(x,\xi,z)+\eta(z)F(x,y,\xi).$$

The intersection of the basic classes is the special class $\mathcal{F}_0$ determined by the condition $F(x,y,z)=0$. Hence $\mathcal{F}_0$ is the class of almost contact B-metric manifolds with ∇-parallel structures, i.e. $\nabla\varphi=\nabla\xi=\nabla\eta=\nabla g=\nabla\tilde{g}=0$.

If $\{e_i;\xi\}$ $(i=1,2,\ldots,2n)$ is a basis of T_pM and $\big(g^{ij}\big)$ is the inverse matrix of (g_{ij}), then with F are associated 1-forms θ, θ^*, ω, called *Lee forms*, defined by:

$$\theta(z)=g^{ij}F(e_i,e_j,z), \quad \theta^*(z)=g^{ij}F(e_i,\varphi e_j,z), \quad \omega(z)=F(\xi,\xi,z).$$

Further, we use the following characteristic conditions of the basic classes [9]:

$$
\begin{aligned}
\mathcal{F}_1: &\quad F(x,y,z) = \tfrac{1}{2n}\big\{g(x,\varphi y)\theta(\varphi z) + g(\varphi x,\varphi y)\theta(\varphi^2 z) \\
&\qquad\qquad + g(x,\varphi z)\theta(\varphi y) + g(\varphi x,\varphi z)\theta(\varphi^2 y)\big\}; \\
\mathcal{F}_2: &\quad F(\xi,y,z) = F(x,\xi,z) = 0, \\
&\quad F(x,y,\varphi z) + F(y,z,\varphi x) + F(z,x,\varphi y) = 0, \quad \theta = 0; \\
\mathcal{F}_3: &\quad F(\xi,y,z) = F(x,\xi,z) = 0, \\
&\quad F(x,y,z) + F(y,z,x) + F(z,x,y) = 0; \\
\mathcal{F}_4: &\quad F(x,y,z) = -\tfrac{1}{2n}\theta(\xi)\big\{g(\varphi x,\varphi y)\eta(z) + g(\varphi x,\varphi z)\eta(y)\big\}; \\
\mathcal{F}_5: &\quad F(x,y,z) = -\tfrac{1}{2n}\theta^*(\xi)\big\{g(x,\varphi y)\eta(z) + g(x,\varphi z)\eta(y)\big\}; \\
\mathcal{F}_6: &\quad F(x,y,z) = F(x,y,\xi)\eta(z) + F(x,z,\xi)\eta(y), \\
&\quad F(x,y,\xi) = F(y,x,\xi) = -F(\varphi x,\varphi y,\xi), \quad \theta = \theta^* = 0; \\
\mathcal{F}_7: &\quad F(x,y,z) = F(x,y,\xi)\eta(z) + F(x,z,\xi)\eta(y), \\
&\quad F(x,y,\xi) = -F(y,x,\xi) = -F(\varphi x,\varphi y,\xi); \\
\mathcal{F}_8: &\quad F(x,y,z) = F(x,y,\xi)\eta(z) + F(x,z,\xi)\eta(y), \\
&\quad F(x,y,\xi) = F(y,x,\xi) = F(\varphi x,\varphi y,\xi); \\
\mathcal{F}_9: &\quad F(x,y,z) = F(x,y,\xi)\eta(z) + F(x,z,\xi)\eta(y), \\
&\quad F(x,y,\xi) = -F(y,x,\xi) = F(\varphi x,\varphi y,\xi); \\
\mathcal{F}_{10}: &\quad F(x,y,z) = F(\xi,\varphi y,\varphi z)\eta(x); \\
\mathcal{F}_{11}: &\quad F(x,y,z) = \eta(x)\left\{\eta(y)\omega(z) + \eta(z)\omega(y)\right\}.
\end{aligned}
\tag{2}
$$

Let now consider the case of the lowest dimension of the manifold (M,φ,ξ,η,g), i.e. $\dim M = 3$.

Let us denote the components $F_{ijk} = F(e_i,e_j,e_k)$ of the tensor F with respect to a *φ-basis* $\{e_i\}_{i=0}^2 = \{e_0 = \xi, e_1 = e, e_2 = \varphi e\}$ of T_pM, which satisfies the following conditions

$$
g(e_0,e_0) = g(e_1,e_1) = -g(e_2,e_2) = 1, \quad g(e_i,e_j) = 0,\ i \neq j \in \{0,1,2\}. \tag{3}
$$

According to [8], the components of the Lee forms with respect to a φ-basis with conditions (3) are

$$
\begin{aligned}
&\theta_0 = F_{110} - F_{220}, && \theta_1 = F_{111} - F_{221}, && \theta_2 = F_{112} - F_{211}, \\
&\theta_0^* = F_{120} + F_{210}, && \theta_1^* = F_{112} + F_{211}, && \theta_2^* = F_{111} + F_{221}, \\
&\omega_0 = 0, && \omega_1 = F_{001}, && \omega_2 = F_{002}.
\end{aligned}
$$

Then, if F_s $(s = 1,2,\dots,11)$ are the components of F in the corresponding basic classes $\mathcal{F}_s$ and $x = x^i e_i$, $y = y^j e_j$, $z = z^k e_k$ are arbitrary vectors we have [8]:

$$
\begin{aligned}
&F_1(x,y,z)=\left(x^1\theta_1-x^2\theta_2\right)\left(y^1z^1+y^2z^2\right),\\
&\quad\theta_1=F_{111}=F_{122},\qquad \theta_2=-F_{211}=-F_{222};\\
&F_2(x,y,z)=F_3(x,y,z)=0;\\
&F_4(x,y,z)=\tfrac{1}{2}\theta_0\Big\{x^1\left(y^0z^1+y^1z^0\right)-x^2\left(y^0z^2+y^2z^0\right)\Big\},\\
&\quad\tfrac{1}{2}\theta_0=F_{101}=F_{110}=-F_{202}=-F_{220};\\
&F_5(x,y,z)=\tfrac{1}{2}\theta_0^*\big\{x^1\left(y^0z^2+y^2z^0\right)+x^2\left(y^0z^1+y^1z^0\right)\big\},\\
&\quad\tfrac{1}{2}\theta_0^*=F_{102}=F_{120}=F_{201}=F_{210};\\
&F_6(x,y,z)=F_7(x,y,z)=0;\\
&F_8(x,y,z)=\lambda\big\{x^1\left(y^0z^1+y^1z^0\right)+x^2\left(y^0z^2+y^2z^0\right)\big\},\\
&\quad\lambda=F_{101}=F_{110}=F_{202}=F_{220};\\
&F_9(x,y,z)=\mu\big\{x^1\left(y^0z^2+y^2z^0\right)-x^2\left(y^0z^1+y^1z^0\right)\big\},\\
&\quad\mu=F_{102}=F_{120}=-F_{201}=-F_{210};\\
&F_{10}(x,y,z)=\nu x^0\left(y^1z^1+y^2z^2\right),\qquad \nu=F_{011}=F_{022};\\
&F_{11}(x,y,z)=x^0\big\{\left(y^1z^0+y^0z^1\right)\omega_1+\left(y^2z^0+y^0z^2\right)\omega_2\big\},\\
&\quad\omega_1=F_{010}=F_{001},\qquad \omega_2=F_{020}=F_{002}.
\end{aligned}
\tag{4}
$$

Obviously, the class of 3-dimensional almost contact B-metric manifolds is

$$
\mathcal{F}_1\oplus\mathcal{F}_4\oplus\mathcal{F}_5\oplus\mathcal{F}_8\oplus\mathcal{F}_9\oplus\mathcal{F}_{10}\oplus\mathcal{F}_{11}. \tag{5}
$$

1.2. **Curvature properties of almost contact B-metric manifolds.** Let $R=[\nabla,\nabla]-\nabla_{[\,,\,]}$ be the curvature (1,3)-tensor of ∇ and the corresponding curvature $(0,4)$-tensor be denoted by the same letter: $R(x,y,z,w)=g(R(x,y)z,w)$. The following properties are valid:

$$
\begin{aligned}
&R(x,y,z,w)=-R(y,x,z,w)=-R(x,y,w,z),\\
&R(x,y,z,w)+R(y,z,x,w)+R(z,x,y,w)=0.
\end{aligned}
\tag{6}
$$

On any $\mathcal{F}_0$-manifold we have

$$
R(x,y,\varphi z,\varphi w)=-R(x,y,z,w). \tag{7}
$$

Any (0,4)-tensor having the properties (6) is called a *curvature-like* tensor. If such a tensor has the Kähler property (7) it is called a *Kähler* tensor.

From [18] it is known that every Kähler tensor on a 3-dimensional almost contact B-metric manifold is zero and therefore every 3-dimensional $\mathcal{F}_0$-manifold is flat, i.e. $R=0$.

The Ricci tensor ρ and the scalar curvature τ for R as well as their associated quantities are defined respectively by

$$
\begin{aligned}
&\rho(y,z)=g^{ij}R(e_i,y,z,e_j), && \tau=g^{ij}\rho(e_i,e_j),\\
&\rho^*(y,z)=g^{ij}R(e_i,y,z,\varphi e_j), && \tau^*=g^{ij}\rho^*(e_i,e_j).
\end{aligned}
$$

Further, we use the notation $g \owedge h$ for the Kulkarni-Nomizu product of two (0,2)-tensors, i.e.

$$\begin{aligned}(g \owedge h)(x,y,z,w) &= g(x,z)h(y,w) - g(y,z)h(x,w)\\ &\quad + g(y,w)h(x,z) - g(x,w)h(y,z).\end{aligned}$$

Obviously, $g \owedge h$ has the properties of R in (6) when g and h are symmetric.

It is well known that the curvature tensor has the following form on any 3-dimensional manifold:

$$R = -g \owedge \left(\rho - \frac{\tau}{4}g\right). \tag{8}$$

Every non-degenerate 2-plane β with respect to g in T_pM, $p \in M$, has the following sectional curvature

$$k(\beta;p) = -\frac{2R(x,y,y,x)}{g \owedge g},$$

where $\{x,y\}$ is a basis of β.

A 2-plane β is said to be a *φ-holomorphic section* (respectively, a *ξ-section*) if $\beta = \varphi\beta$ (respectively, $\xi \in \beta$).

2. Lie Groups as 3-Dimensional Almost Contact B-Metric Manifolds

Let L be a 3-dimensional real connected Lie group, and $\mathfrak{l}$ be its Lie algebra with a basis $\{E_0, E_1, E_2\}$ of left invariant vector fields. Let an almost contact structure (φ, ξ, η) be defined by

$$\begin{aligned}&\varphi E_0 = 0, \quad \varphi E_1 = E_2, \quad \varphi E_2 = -E_1, \quad \xi = E_0,\\ &\eta(E_0) = 1, \quad \eta(E_1) = \eta(E_2) = 0\end{aligned} \tag{9}$$

and let g be a pseudo-Riemannian metric such that

$$\begin{aligned}&g(E_0,E_0) = g(E_1,E_1) = -g(E_2,E_2) = 1,\\ &g(E_0,E_1) = g(E_0,E_2) = g(E_1,E_2) = 0.\end{aligned} \tag{10}$$

Then we have a 3-dimensional almost contact B-metric manifold $(L, \varphi, \xi, \eta, g)$.

The corresponding Lie algebra $\mathfrak{l}$ is determined as follows:

$$[E_i, E_j] = C_{ij}^k E_k, \quad i,j,k \in \{0,1,2\}. \tag{11}$$

Applying the Jacobi identity

$$\big[[E_i,E_j],E_k\big] + \big[[E_j,E_k],E_i\big] + \big[[E_k,E_i],E_j\big] = 0,$$

from the nine commutation coefficients C_{ij}^k remain six which could be chosen as parameters, so the three coefficients with different indices are expressed by the six parameters (if the denominators are non-zero) as follows:

$$C_{12}^0 = \frac{C_{12}^1 C_{01}^0 + C_{12}^2 C_{02}^0}{C_{01}^1 + C_{02}^2}, \qquad C_{02}^1 = \frac{C_{01}^1 C_{02}^0 - C_{12}^1 C_{02}^2}{C_{01}^0 - C_{12}^2},$$
$$C_{01}^2 = \frac{C_{01}^0 C_{02}^2 + C_{01}^1 C_{12}^2}{C_{02}^0 + C_{12}^1}.$$

The Koszul equality

$$2g\left(\nabla_{E_i}E_j,E_k\right)=g\left([E_i,E_j],E_k\right)+g\left([E_k,E_i],E_j\right)+g\left([E_k,E_j],E_i\right) \tag{12}$$

implies the following formula for the components $F_{ijk}=F(E_i,E_j,E_k)$, $i,j,k\in\{0,1,2\}$, of the tensor F:

$$\begin{aligned}2F_{ijk}=g\left([E_i,\varphi E_j]-\varphi[E_i,E_j],E_k\right)+g\left(\varphi[E_k,E_i]-[\varphi E_k,E_i],E_j\right)\\ +g\left([E_k,\varphi E_j]-[\varphi E_k,E_j],E_i\right).\end{aligned}$$

Then we have

$$\begin{aligned}&F_{111}=F_{122}=2C^1_{12}, && F_{211}=F_{222}=2C^2_{12},\\ &F_{120}=F_{102}=-C^1_{01}, && F_{020}=F_{002}=-C^0_{01},\\ &F_{210}=F_{201}=-C^2_{02}, && F_{010}=F_{001}=C^0_{02},\\ &F_{110}=F_{101}=\tfrac12\left(C^0_{12}-C^2_{01}+C^1_{02}\right), &&\\ &F_{220}=F_{202}=\tfrac12\left(C^0_{12}+C^2_{01}-C^1_{02}\right), &&\\ &F_{011}=F_{022}=C^0_{12}+C^2_{01}+C^1_{02}. &&\end{aligned} \tag{13}$$

For the Lee forms we obtain

$$\begin{aligned}&\theta_0=-C^2_{01}+C^1_{02}, && \theta^*_0=-C^1_{01}-C^2_{02}, && \omega_0=0,\\ &\theta_1=2C^1_{12}, && \theta^*_1=2C^2_{12}, && \omega_1=C^0_{02},\\ &\theta_2=-2C^2_{12}, && \theta^*_2=2C^1_{12}, && \omega_2=-C^0_{01}.\end{aligned} \tag{14}$$

Using (4), (13) and (14), we deduce the following

Theorem 2.1. *The manifold (L,φ,ξ,η,g) belongs to the basic class $\mathcal{F}_s$ ($s\in\{1,4,5,8,9,10,11\}$) if and only if the corresponding Lie algebra $\mathfrak{l}$ is determined by the following commutators:*

$$\begin{aligned}&\mathcal{F}_1: && [E_0,E_1]=[E_0,E_2]=0, && [E_1,E_2]=\alpha E_1+\beta E_2;\\ &\mathcal{F}_4: && [E_0,E_1]=\alpha E_2, && [E_0,E_2]=-\alpha E_1, && [E_1,E_2]=0;\\ &\mathcal{F}_5: && [E_0,E_1]=\alpha E_1, && [E_0,E_2]=\alpha E_2, && [E_1,E_2]=0;\\ &\mathcal{F}_8: && [E_0,E_1]=\alpha E_2, && [E_0,E_2]=\alpha E_1, && [E_1,E_2]=-2\alpha E_0;\\ &\mathcal{F}_9: && [E_0,E_1]=\alpha E_1, && [E_0,E_2]=-\alpha E_2, && [E_1,E_2]=0;\\ &\mathcal{F}_{10}: && [E_0,E_1]=\alpha E_2, && [E_0,E_2]=\alpha E_1, && [E_1,E_2]=0;\\ &\mathcal{F}_{11}: && [E_0,E_1]=\alpha E_0, && [E_0,E_2]=\beta E_0, && [E_1,E_2]=0,\end{aligned} \tag{15}$$

where α, β are arbitrary real parameters. Moreover, the relations of α and β with the non-zero components F_{ijk} in the different basic classes $\mathcal{F}_s$ from (4) *are the following:*

$$\begin{aligned}&\mathcal{F}_1: && \alpha=\tfrac12\theta_1, \quad \beta=\tfrac12\theta_2; && \mathcal{F}_4: && \alpha=\tfrac12\theta_0;\\ &\mathcal{F}_5: && \alpha=-\tfrac12\theta^*_0; && \mathcal{F}_8: && \alpha=-\lambda;\\ &\mathcal{F}_9: && \alpha=-\mu; && \mathcal{F}_{10}: && \alpha=\tfrac12\nu;\\ &\mathcal{F}_{11}: && \alpha=-\omega_2, \quad \beta=\omega_1. &&&&\end{aligned}$$

Let us remark that an $\mathcal{F}_0$-manifold is obtained when the Lie algebra is Abelian, i.e. all commutators are zero. Further, we omit this special case.

2.1. **Structures of special type on the considered manifolds.** The metric g is called *Killing* (or, Ad-*invariant*) if

$$g([x,y],z) = g(x,[y,z]). \tag{16}$$

Bearing in mind (10), (11) and (16), we obtain that g is Killing if and only if the equalities $C^0_{12} = -C^2_{01} = -C^1_{02}$ are valid and the rest of the commutation coefficients are zero. Then, according to (13) and (14), we get the following non-zero components $-F_{011} = -F_{022} = 2F_{110} = 2F_{101} = 2F_{220} = 2F_{202} = C^0_{12}$ and $\theta = \theta^* = \omega = 0$. Therefore, using (4), we establish that $F = F_8 + F_{10}$ and $2\lambda = -\nu$. Thus, it is valid the following

Theorem 2.2. *The metric g is Killing if and only if (L,φ,ξ,η,g) belongs to the subclass of $\mathcal{F}_8 \oplus \mathcal{F}_{10}$ determined by the condition $2\lambda = -\nu$.*

In a similar way we obtain that $\tilde{g}$ is Killing, i.e. $\tilde{g}([x,y],z) = \tilde{g}(x,[y,z])$, if and only if $C^0_{12} = -C^1_{01} = C^2_{02}$ and the others C^k_{ij} are zero. Then from (13) we have zero Lee forms and $F_{120} = F_{102} = -F_{210} = -F_{201} = 2F_{110} = 2F_{101} = 2F_{220} = 2F_{202} = F_{011} = F_{022} = C^0_{12}$, i.e. $2\lambda = \mu = \nu$, bearing in mind (4). Therefore, we get

Theorem 2.3. *The metric $\tilde{g}$ is Killing if and only if (L,φ,ξ,η,g) belongs to the subclass of $\mathcal{F}_8 \oplus \mathcal{F}_9 \oplus \mathcal{F}_{10}$ determined by the condition $2\lambda = \mu = \nu$.*

The structure φ is called *bi-invariant*, if $\varphi[x,y] = [x,\varphi y]$. This condition is satisfied if and only if $F_{011} = F_{022} = 2F_{110} = 2F_{101} = 2F_{220} = 2F_{202} = C^0_{12}$, i.e. $2\lambda = \nu$. Therefore, it is valid the following

Theorem 2.4. *The structure φ is bi-invariant if and only if (L,φ,ξ,η,g) belongs to the subclass of $\mathcal{F}_8 \oplus \mathcal{F}_{10}$ determined by the condition $2\lambda = \nu$.*

The structure φ is called *Abelian*, if $[\varphi x, \varphi y] = [x,y]$. This condition is valid if and only if $F_{011} = F_{022} = 2F_{110} = 2F_{101} = 2F_{220} = 2F_{202} = C^0_{12}$, i.e. $2\lambda = \nu$, and $F_{111} = F_{122} = 2C^1_{12} = \theta_1$, $F_{211} = F_{222} = 2C^2_{12} = -\theta_2$. Therefore, we deduce

Theorem 2.5. *The structure φ is Abelian if and only if (L,φ,ξ,η,g) belongs to the subclass of $\mathcal{F}_1 \oplus \mathcal{F}_8 \oplus \mathcal{F}_{10}$ determined by the condition $2\lambda = \nu$.*

The vector field ξ is called *Killing*, if the Lie derivative with respect to ξ of the metric g is zero, i.e. $\mathcal{L}_\xi g = 0$. Then the manifold (L,φ,ξ,η,g) is called a *K-contact B-metric manifold*. This condition is equivalent to $(\nabla_x \eta)\, y + (\nabla_y \eta)\, x = 0$, which is satisfied for any manifold in $\mathcal{F}_1 \oplus \mathcal{F}_2 \oplus \mathcal{F}_3 \oplus \mathcal{F}_7 \oplus \mathcal{F}_8 \oplus \mathcal{F}_{10}$ for arbitrary dimension, according to (2). Then, bearing in mind (5) – the class of the considered manifolds of dimension 3, we get

Theorem 2.6. *The vector field ξ is Killing if and only if $(L,\varphi,\xi,\eta,g) \in \mathcal{F}_1 \oplus \mathcal{F}_8 \oplus \mathcal{F}_{10}$, i.e. the class of the 3-dimensional K-contact B-metric manifolds is $\mathcal{F}_1 \oplus \mathcal{F}_8 \oplus \mathcal{F}_{10}$.*

2.2. **Curvature properties of the considered manifolds.** Using (12) and (15), we obtain the covariant derivatives of E_i with respect to ∇. The non-zero ones of

them are the following:

$$\begin{aligned}
\mathcal{F}_1: &\quad \nabla_{E_1}E_1=\alpha E_2,\ \nabla_{E_1}E_2=\alpha E_1,\ \nabla_{E_2}E_1=-\beta E_2,\ \nabla_{E_2}E_2=-\beta E_1;\\
\mathcal{F}_4: &\quad \nabla_{E_1}E_0=-\alpha E_2,\ \nabla_{E_2}E_0=\alpha E_1;\\
\mathcal{F}_5: &\quad \nabla_{E_1}E_1=-\nabla_{E_2}E_2=\alpha E_0,\ \nabla_{E_1}E_0=-\alpha E_1,\ \nabla_{E_2}E_0=-\alpha E_2;\\
\mathcal{F}_8: &\quad \nabla_{E_1}E_2=-\nabla_{E_2}E_1=-\alpha E_0,\ \nabla_{E_1}E_0=-\alpha E_2,\ \nabla_{E_2}E_0=-\alpha E_1;\\
\mathcal{F}_9: &\quad \nabla_{E_1}E_1=\nabla_{E_2}E_2=\alpha E_0,\ \nabla_{E_1}E_0=-\alpha E_1,\ \nabla_{E_2}E_0=\alpha E_2;\\
\mathcal{F}_{10}: &\quad \nabla_{E_0}E_1=\alpha E_2,\ \nabla_{E_0}E_2=\alpha E_1;\\
\mathcal{F}_{11}: &\quad \nabla_{E_0}E_0=-\alpha E_1+\beta E_2,\ \nabla_{E_0}E_1=\alpha E_0,\nabla_{E_0}E_2=\beta E_0.
\end{aligned}$$

By virtue of the latter equalities and bearing in mind the definition equalities of the curvature tensor R, the Ricci tensor ρ, the $*$-Ricci tensor ρ^*, the scalar curvature τ, its $*$-scalar curvature τ^* and the sectional curvature k_{ij} of the non-degenerate 2-plane $\{E_i,E_j\}$

$$\begin{aligned}
&R_{ijkl}=g\left(\nabla_{E_i}\nabla_{E_j}E_k-\nabla_{E_j}\nabla_{E_i}E_k-\nabla_{[E_i,E_j]}E_k,E_l\right),\\
&\rho_{jk}=R_{0jk0}+R_{1jk1}-R_{2jk2},\qquad \rho^*_{jk}=R_{1jk2}+R_{2jk1},\\
&\tau=\rho_{00}+\rho_{11}-\rho_{22},\qquad \tau^*=2\rho_{12},\qquad k_{ij}=-2R_{ijij}/(g\owedge g)_{ijij},
\end{aligned}\tag{17}$$

from Theorem 2.1 we obtain the following

Theorem 2.7. *The $\mathcal{F}_{10}$-manifolds (L,φ,ξ,η,g) are flat and the non-zero components of R, ρ, ρ^* and the non-zero values of τ, τ^*, k_{ij} for the manifolds (L,φ,ξ,η,g) from the rest of the basic classes are the following:*

$$\begin{aligned}
\mathcal{F}_1: &\quad R_{1212}=\rho_{11}=-\rho_{22}=\rho^*_{12}=\rho^*_{21}=\tfrac{1}{2}\tau=k_{12}=\alpha^2-\beta^2;\\
\mathcal{F}_4: &\quad -R_{0101}=R_{0202}=\tfrac{1}{2}\rho_{00}=\rho_{11}=-\rho_{22}\\
&\qquad\qquad =\tfrac{1}{4}\tau=k_{01}=k_{02}=\alpha^2;\\
\mathcal{F}_5: &\quad -R_{0101}=R_{0202}=R_{1212}=\tfrac{1}{2}\rho_{00}=\tfrac{1}{2}\rho_{11}=-\tfrac{1}{2}\rho_{22}\\
&\qquad\qquad =\rho^*_{12}=\rho^*_{21}=\tfrac{1}{6}\tau=k_{12}=k_{01}=k_{02}=-\alpha^2;\\
\mathcal{F}_8: &\quad R_{0101}=-R_{0202}=R_{1212}=-\tfrac{1}{2}\rho_{00}=\rho^*_{12}=\rho^*_{21}\\
&\qquad\qquad =-\tfrac{1}{2}\tau=k_{12}=-k_{01}=-k_{02}=\alpha^2;\\
\mathcal{F}_9: &\quad R_{0101}=-R_{0202}=R_{1212}=-\tfrac{1}{2}\rho_{00}=\rho^*_{12}=\rho^*_{21}\\
&\qquad\qquad =-\tfrac{1}{2}\tau=k_{12}=-k_{01}=-k_{02}=\alpha^2;\\
\mathcal{F}_{11}: &\quad R_{0101}=-\rho_{11}=-k_{01}=\alpha^2,\quad R_{0202}=-\rho_{22}=k_{02}=\beta^2,\\
&\quad R_{0120}=\rho_{12}=\tfrac{1}{2}\tau^*=-\alpha\beta,\quad \rho_{00}=\tfrac{1}{2}\tau=-\alpha^2+\beta^2.
\end{aligned}\tag{18}$$

Let us remark, on the 3-dimensional almost contact B-metric manifold with the basis $\{E_0,E_1,E_2\}$ defined by (9), we have two basic ξ-sections $\{E_0,E_1\}$, $\{E_0,E_2\}$ and one basic φ-holomorphic section $\{E_1,E_2\}$.

According to Theorem 2.7, we obtain

Theorem 2.8. *For the $\mathcal{F}_s$-manifolds (L,φ,ξ,η,g), which do not belong to $\mathcal{F}_0$, the following propositions are valid:*

(1) *Only the $\mathcal{F}_1$-manifolds have a Kähler tensor R;*

(2) *An $\mathcal{F}_1$-manifold is flat if and only if $\alpha^2 = \beta^2$;*
(3) *All $\mathcal{F}_8$-manifolds and $\mathcal{F}_9$-manifolds have curvature tensors of the same form;*
(4) *The curvature tensor of any $\mathcal{F}_s$-manifold ($s = 4, 11$) has the property $R(x, y, \varphi z, \varphi w) = 0$;*
(5) *Every $\mathcal{F}_s$-manifold ($s = 4, 11$) has a vanishing $*$-Ricci tensor;*
(6) *Every $\mathcal{F}_s$-manifold ($s = 4, 8, 9$) has a positive scalar curvature;*
(7) *An $\mathcal{F}_1$-manifold (resp., $\mathcal{F}_{11}$-manifold) has a positive scalar curvature if and only if $\alpha^2 > \beta^2$ (resp., $\alpha^2 < \beta^2$);*
(8) *Every $\mathcal{F}_5$-manifold has a negative scalar curvature;*
(9) *An $\mathcal{F}_1$-manifold (resp., $\mathcal{F}_{11}$-manifold) has a negative scalar curvature if and only if $\alpha^2 < \beta^2$ (resp., $\alpha^2 > \beta^2$);*
(10) *Every $\mathcal{F}_s$-manifold ($s = 1, 4, 5, 8, 9$) has a vanishing $*$-scalar curvature;*
(11) *An $\mathcal{F}_{11}$-manifold has a vanishing $*$-scalar curvature if and only if $\alpha\beta = 0$ for $\alpha \neq \beta$;*
(12) *An $\mathcal{F}_{11}$-manifold has a positive (resp., negative) $*$-scalar curvature if and only if $\alpha\beta < 0$ (resp., $\alpha\beta > 0$);*
(13) *Every $\mathcal{F}_1$-manifold has vanishing sectional curvatures of the ξ-sections;*
(14) *Every $\mathcal{F}_4$-manifold has positive sectional curvatures of the basic ξ-sections;*
(15) *Every $\mathcal{F}_s$-manifold ($s = 5, 8, 9$) has negative sectional curvatures of the basic ξ-sections;*
(16) *An $\mathcal{F}_1$-manifold has a vanishing scalar curvature of the basic φ-holomorphic section if and only if the manifold is flat;*
(17) *Every $\mathcal{F}_s$-manifold ($s = 4, 11$) has a vanishing scalar curvature of the basic φ-holomorphic section;*
(18) *Every $\mathcal{F}_s$-manifold ($s = 8, 9$) (resp., $\mathcal{F}_5$-manifold) has a positive (resp., negative) scalar curvature of the basic φ-holomorphic section;*
(19) *An $\mathcal{F}_1$-manifold has a positive (resp., negative) scalar curvature of the basic φ-holomorphic section if and only if $\alpha^2 > \beta^2$ (resp., $\alpha^2 < \beta^2$).*

Taking into account (18), we get the following

Corollary 2.9. *The form of the Ricci tensor on $(L, \varphi, \xi, \eta, g)$ in the respective basic class is:*

$$\begin{aligned}
&\mathcal{F}_1: \quad \rho = \tfrac{\tau}{2}(g - \eta\otimes\eta); && \mathcal{F}_4: \quad \rho = \tfrac{\tau}{4}(g + \eta\otimes\eta);\\
&\mathcal{F}_5: \quad \rho = \tfrac{\tau}{3}g; && \mathcal{F}_8: \quad \rho = \tau(\eta\otimes\eta);\\
&\mathcal{F}_9: \quad \rho = \tau(\eta\otimes\eta); && \mathcal{F}_{11}: \quad \rho = \rho(\varphi\cdot,\varphi\cdot) + \tfrac{\tau}{2}g - \tau^* g^*,
\end{aligned}$$

where $g^*(\cdot,\cdot) = g(\cdot,\varphi\cdot) = \tilde{g} - \eta\otimes\eta$.

Using Corollary 2.9 and (8), we obtain for the following

Corollary 2.10. *The form of the curvature tensor on (L,φ,ξ,η,g) in the respective basic class is:*

$$\begin{aligned}
\mathcal{F}_1:&\quad R=-\tfrac{\tau}{4}\,(g\owedge g)+\tfrac{\tau}{2}\,(g\owedge(\eta\otimes\eta))\,;\\
\mathcal{F}_4:&\quad R=-\tfrac{\tau}{4}\,(g\owedge(\eta\otimes\eta))\,;\\
\mathcal{F}_5:&\quad R=-\tfrac{\tau}{12}\,(g^*\owedge g^*)-\tfrac{\tau}{6}\,(g\owedge(\eta\otimes\eta))\,;\\
\mathcal{F}_8:&\quad R=\tfrac{\tau}{4}\,(g^*\owedge g^*)-\tfrac{\tau}{2}\,(g\owedge(\eta\otimes\eta))\,;\\
\mathcal{F}_9:&\quad R=\tfrac{\tau}{4}\,(g^*\owedge g^*)-\tfrac{\tau}{2}\,(g\owedge(\eta\otimes\eta))\,;\\
\mathcal{F}_{11}:&\quad R=-\rho\owedge(\eta\otimes\eta).
\end{aligned}$$

Let (V,φ,ξ,η,g) be a vector space endowed with an almost contact structure (φ,ξ,η). The decomposition $x=-\varphi^2x+\eta(x)\xi$ generates the projectors h and v on (V,φ,ξ,η) determined by $h(x)=-\varphi^2x$ and $v(x)=\eta(x)\xi$ and having the properties $h\circ h=h$, $v\circ v=v$, $h\circ v=v\circ h=0$. Therefore, we have the orthogonal decomposition $V=h(V)\oplus v(V)$. Then we obtain the corresponding linear operators ℓ_1,ℓ_2,ℓ_3 in the space $V^*\times V^*$ of the tensors S of type (0,2) over (V,φ,ξ,η) defined in [1]:

$$\ell_1(S)(x,y)=S(h(x),h(y)),\quad \ell_2(S)(x,y)=S(v(x),v(y)),$$
$$\ell_3(S)(x,y)=S(v(x),y)+S(x,v(y))-2S(v(x),v(y)).$$

The structure group $\mathcal{G}$ of (M,φ,ξ,η,g) is determined by $\mathcal{G}=\mathcal{O}(n;\mathbb{C})\times\mathcal{I}$, where $\mathcal{I}$ is the identity on $\mathrm{Span}(\xi)$ and $\mathcal{O}(n;\mathbb{C})=\mathcal{GL}(n;\mathbb{C})\cap\mathcal{O}(n,n)$.

It is known [10], the following orthogonal decomposition which is invariant with respect to the action of the structure group $\mathcal{G}$:

$$V^*\times V^*=\ell_1(V^*\times V^*)\oplus\ell_2(V^*\times V^*)\oplus\ell_3(V^*\times V^*),$$

where

$$\ell_i(V^*\times V^*)=\{S\in(V^*\times V^*)\mid S=\ell_i(S)\},\quad i=1,2,3.$$

Since the metrics g and $\tilde g$ belong to $V^*\times V^*$, then their components in the three orthogonal subspaces are:

$$\begin{aligned}
&\ell_1(g)=-g(\varphi\cdot,\varphi\cdot)=g-\eta\otimes\eta, && \ell_2(g)=\eta\otimes\eta, && \ell_3(g)=0,\\
&\ell_1(\tilde g)=g(\cdot,\varphi\cdot)=\tilde g-\eta\otimes\eta, && \ell_2(\tilde g)=\eta\otimes\eta, && \ell_3(\tilde g)=0.
\end{aligned}$$

On any almost contact B-metric manifold there are a pair of B-metrics $(g,\tilde g)$ which restrictions on $H=\mathrm{Ker}(\eta)$ coincide with their ℓ_1-images and are the real part and the imaginary part of the complex Riemannian metric $g^{\mathbb{C}}$, i.e. $g^{\mathbb{C}}=g|_H+i\tilde g|_H=\ell_1(g)+i\ell_1(\tilde g)$. Moreover, the restrictions of g and $\tilde g$ on $H^\perp=\mathrm{Span}(\xi)$ coincide with their ℓ_2-images and $\eta\otimes\eta$, i.e. $g|_{H^\perp}=\tilde g|_{H^\perp}=\ell_2(g)=\ell_2(\tilde g)=\eta\otimes\eta$.

It is well known that a manifold is called *Einstein* if the Ricci tensor is proportional to the metric tensor, i.e. $\rho=\lambda g$, $\lambda\in\mathbb{R}$.

Then, bearing in mind the above arguments, it is reasonable to call a manifold M whose Ricci tensor satisfies the condition

$$\rho=\lambda g+\mu\tilde g+\nu\eta\otimes\eta \tag{19}$$

an *η-complex-Einstein manifold*. The case of the η-complex-Einstein Sasaki-like manifolds is considered in [7]. If $\mu=0$ we call M an *η-Einstein manifold*. On

the almost contact metric manifolds in [3] is studied the η-Einstein geometry as a class of distinguished Riemannian metrics on the contact metric manifolds. It is well-known [22] that λ and ν are constants if the almost contact metric manifold is K-contact and has dimension greater than 3.

It is meaningful to consider the Einstein property in the separate components $\ell_i(\rho)$ and $\ell_i(\rho^*)$, $i = 1, 2, 3$.

In [18] the following notions related to the Einstein condition are introduced. An almost contact B-metric manifold $(M, \varphi, \xi, \eta, g)$ is called *contact-Einstein* if the Ricci tensor has the form $\rho = \lambda g|_H + \mu\tilde{g}|_H + \nu\eta \otimes \eta$, where $\lambda, \mu, \nu \in \mathbb{R}$. A contact-Einstein manifold is called *h-Einstein* (resp., *v-Einstein*) if $\rho = \lambda g|_H + \mu\tilde{g}|_H$ (resp., $\rho = \nu\eta \otimes \eta$). An h-Einstein manifold is called *φ-Einstein* (resp., *$*$-Einstein*) if $\rho = \lambda g|_H$ (resp., $\rho = \mu\tilde{g}|_H$).

According to Corollary 2.9, we obtain

Proposition 2.11. *The manifold $(M, \varphi, \xi, \eta, g)$ is:*

(1) *φ-Einstein if it belongs to $\mathcal{F}_1$;*
(2) *η-Einstein if it belongs to $\mathcal{F}_4$;*
(3) *Einstein if it belongs to $\mathcal{F}_5$;*
(4) *v-Einstein if it belongs to $\mathcal{F}_8 \oplus \mathcal{F}_9$.*

3. An Example

In [8] is introduced a real connected Lie group L as a manifold from the class $\mathcal{F}_9 \oplus \mathcal{F}_{10}$. We consider now this example for $\dim L = 3$ and give geometrical characteristics in relation with some previous results in this work. In this case L is a 3-dimensional real connected Lie group and its associated Lie algebra with a global basis $\{E_0, E_1, E_2\}$ of left invariant vector fields on L is defined by

$$[E_0, E_1] = -a_1E_1 - a_2E_2, \quad [E_0, E_2] = -a_2E_1 + a_1E_2, \quad [E_1, E_2] = 0,$$

where a_1, a_2 are real constants. Actually, L is a family of Lie groups depending on the pair of real parameters. An invariant almost contact structure is defined by (9) and g is a pseudo-Riemannian metric such that (10) are valid. Thus, because of (1), the induced 3-dimensional manifold $(L, \varphi, \xi, \eta, g)$ is an almost contact B-metric manifold.

Let us remark that in [21] the same Lie group with the same almost contact structure but equipped with a compatible Riemannian metric is studied as an almost cosymplectic manifold.

The non-zero components F_{ijk} of the tensor F are following:

$$F_{011} = F_{022} = -2a_2, \quad F_{102} = F_{120} = -F_{201} = -F_{210} = a_1.$$

Thus, it follows that $(L, \varphi, \xi, \eta, g) \in \mathcal{F}_9 \oplus \mathcal{F}_{10}$, bearing in mind (4) for $\mu = a_1$, $\nu = -2a_2$.

Further, we continue with examination of this family of almost contact B-metric manifolds.

Using (12), we obtain the covariant derivatives with respect to ∇ of E_i for the manifold (L,φ,ξ,η,g):

$$\nabla_{E_0}E_1=\tfrac{\nu}{2}E_2,\quad \nabla_{E_0}E_2=\tfrac{\nu}{2}E_1,\quad \nabla_{E_1}E_0=\mu E_1,\quad \nabla_{E_2}E_0=-\mu E_2,$$
$$\nabla_{E_1}E_1=\nabla_{E_2}E_2=-\mu E_0.$$

By virtue of the latter equalities and bearing in mind the definition equalities (17), we get:

$$\begin{gathered}R_{0110}=-R_{0220}=R_{1221}=-\mu^2,\qquad R_{0120}=\mu\nu,\\ \rho_{00}=-2\rho_{22}=-2\mu^2,\qquad \rho^*_{00}=2\rho_{12}=2\mu\nu,\\ \tau=-2\mu^2,\qquad \tau^*=2\mu\nu,\\ k_{01}=k_{02}=-\mu^2,\qquad k_{12}=\mu^2\end{gathered}\tag{20}$$

We obtain the form of the Ricci tensor ρ:

$$\rho=\left(\tau+\frac{\tau^*}{2}\right)\eta\otimes\eta-\frac{\tau^*}{2}\tilde{g}.\tag{21}$$

Consequently, bearing in mind (19), (L,φ,ξ,η,g) is an η-complex-Einstein manifold with $\lambda=0$.

According to (8) and (21), we obtain the following form of the curvature tensor:

$$R=\frac{1}{4}g\owedge\Big(\tau g+2\tau^*\tilde{g}-2\left(2\tau+\tau^*\right)\eta\otimes\eta\Big).\tag{22}$$

If $\mu=0$, $\nu\neq0$, i.e. the manifold belongs to $\mathcal{F}_{10}$, then $\tau=\tau^*=0$. Therefore, (22) implies $R=0$.

If $\nu=0$, $\mu\neq0$, i.e. the manifold belongs to $\mathcal{F}_9$, then $\tau^*=0$, $\tau\neq0$ and

$$R=\frac{\tau}{4}g\owedge\Big(g-4\eta\otimes\eta\Big).$$

The latter conclusions support Corollary 2.9 and Corollary 2.10.

Taking into account (20), we obtain that the sectional curvatures of the basic ξ-sections (respectively, the basic φ-holomorphic sections) are negative (respectively, positive) for $(L,\varphi,\xi,\eta,g)\in\mathcal{F}_9\oplus\mathcal{F}_{10}$, which support Theorem 2.8 for $\mathcal{F}_9$ and $\mathcal{F}_{10}$.

Acknowledgment. This paper is partially supported by project NI13-FMI-002 of the Scientific Research Fund, Plovdiv University, Bulgaria

References

[1] Alexiev, V., Ganchev, G.: *On the classification of almost contact metric manifolds*. In: Math. and Educ. in Math. Proc. 15th Spring Conf. of UBM. Sunny Beach, 155–161 (1986)

[2] Blair, D.E.: *Riemannian Geometry of Contact and Symplectic Manifolds*. Progress in Mathematics **203**. Birkhäuser, Boston (2002)

[3] Boyer, C.P., Galicki, K., Matzeu, P.: *On η-Einstein Sasakian geometry*. Commun. Math. Phys. **262**, 177–208 (2006)

[4] Ganchev, G., Borisov, A.: *Note on the almost complex manifolds with a Norden metric*. C. R. Acad. Bulg. Sci. **39**, 31–34 (1986)

[5] Ganchev, G., Mihova, V., Gribachev, K.: *Almost contact manifolds with B-metric*. Math. Balkanica (N.S.) **7** (3-4), 261–276 (1993)

[6] Gribachev, K., Mekerov, D., Djelepov, G.: *Generalized B-manifolds*. C. R. Acad. Bulg. Sci. **38**, 299–302 (1985)
[7] Ivanov, S., Manev, H., Manev, M.: *Sasaki-like almost contact complex Riemannian manifolds*. arXiv:1402.5426
[8] Manev, H.: *On the structure tensors of almost contact B-metric manifolds*. arXiv:1405.3088
[9] Manev, M.: *Almost contact B-metric hypersurfaces of Kaehlerian manifolds with B-metric*. In: Dimiev, S., Sekigawa, K. (eds.) Perspectives of Complex Analysis, Differential Geometry and Mathematical Physics, pp. 159–170. World Sci. Publ., Singapore (2001)
[10] Manev, M.: *On the conformal geometry of almost contact manifolds with B-metric*. Ph. D. Thesis. University of Plovdiv, (1998)
[11] Manev, M.: *Curvature properties on some classes of almost contact manifolds with B-metric*. C. R. Acad. Bulg. Sci. **65** (3), 283–290 (2012)
[12] Manev, M.: *Contactly conformal transformations of general type of almost contact manifolds with B-metric. Applications*. Math. Balkanica (N.S.) **11** (3-4), 347–357 (1997)
[13] Manev, M.: *Natural connection with totally skew-symmetric torsion on almost contact manifolds with B-metric*. Int. J. Geom. Methods Mod. Phys. **9** (5), 1250044 (20 pages) (2012)
[14] Manev, M., Gribachev, K.: *Contactly conformal transformations of almost contact manifolds with B-metric*. Serdica Math. J. **19**, 287–299 (1993)
[15] Manev, M., Gribachev, K.: *Conformally invariant tensors on almost contact manifolds with B-metric*. Serdica Math. J. **20**, 133–147 (1994)
[16] Manev, M., Ivanova, M.: *Canonical type connections on almost contact manifold with B-matric*. Ann. Global Anal. Geom. **43** (4), 397–408 (2013)
[17] Manev, M., Ivanova, M.: *A classification of the torsion tensors on almost contact manifolds with B-metric*. Cent. Eur. J. Math. **12** (10), 1416–1432 (2014)
[18] Manev, M., Nakova, G.: *Curvature properties of some three-dimensional almost contact B-metric manifolds*. Plovdiv Univ. Sci. Works – Math. **34** (3), 51–60 (2004)
[19] Nakova, G., Gribachev, K.: *Submanifolds of some almost contact manifolds with B-metric with codimension two, I*. Math. Balkanica (N.S.) **11**, 255–267 (1997)
[20] Manev, M., Nakova, G.: *Curvature properties on some three-dimensional almost contact manifolds with B-metric, II*. In: Mladenov, I.M., Hirshfeld, A.C. (eds.) Proc. 5th Internat. Conf. Geometry, Integrability & Quantization, V, pp. 169–177. SOFTEX, Sofia (2004)
[21] Olszak, Z.: *On almost cosymplectic manifolds*. Kodai Math. J. **4** (2), 239–250 (1981)
[22] Yano, K., Kon, M.: *Structures on Manifolds*. Series in Pure Mathematics **3**, World Sci. Publ., Singapore (1984)

(H. Manev) Medical University of Plovdiv, Faculty of Pharmacy, Department of Pharmaceutical Sciences, 15-A Vasil Aprilov Blvd., Plovdiv 4002, Bulgaria;

Paisii Hilendarski University of Plovdiv, Faculty of Mathematics and Informatics, Department of Algebra and Geometry, 236 Bulgaria Blvd., Plovdiv 4027, Bulgaria

E-mail address: hmanev@uni-plovdiv.bg

(D. Mekerov) Paisii Hilendarski University of Plovdiv, Faculty of Mathematics and Informatics, Department of Algebra and Geometry, 236 Bulgaria Blvd., Plovdiv 4027, Bulgaria

E-mail address: mircho@uni-plovdiv.bg